

**Association between passion and optimal functioning in autistic individuals: the Dualistic Model for Passion**

Alexa Meilleur<sup>1</sup>, Noémie Cusson<sup>1</sup>, Robert J. Vallerand<sup>1</sup>, Mélanie Couture<sup>3</sup>, Elsa Gilbert<sup>4</sup>,  
Isabelle Soulières<sup>1</sup>, Eve-Line Bussières<sup>2</sup>

**Affiliations**

- 1) Psychology department, Université du Québec à Montréal, Montréal, QC, Canada
- 2) Psychology department, Université du Québec à Trois-Rivières, Trois-Rivières, QC, Canada
- 3) School of Rehabilitation, Sherbrooke University, Sherbrooke, QC, Canada
- 4) Health Sciences, Université du Québec à Rimouski, Rimouski, QC, Canada

Corresponding author: Alexa Meilleur, [Meilleur.alex@courrier.uqam.ca](mailto:Meilleur.alex@courrier.uqam.ca)

**Autism in Adulthood, in press**

**Abstract**

**Background:** Autistic individuals have intense interests in which they invest a significant amount of time and energy. Intense interests (i.e., passions) and their impact on optimal functioning were investigated for the first time using the Dualistic Model of Passion (DMP). The DMP posits that harmonious (HP) and obsessive passions (OP) can predict optimal functioning (well-being, contribution to society, and performance). Whereas HP is described as a balanced and flexible form of engagement towards a topic or activity, OP is defined as a rigid form of engagement which negatively impacts optimal functioning.

**Methods:** Autistic individuals aged 14 to 33 ( $n = 108$ ) participated in an online quantitative study and completed self-report measures relating to their favorite interest of the moment (i.e., HP, OP, emotions, flow, conflict, rumination, and optimal functioning). Aims were to characterize passion and to determine whether HP and OP predicted emotions, flow, conflict, rumination, and optimal functioning. Descriptive analyses (means, standard deviations, and pairwise correlations) and a path analysis model were performed to answer these aims.

**Results:** Results revealed that participants were highly passionate for their favorite interest, showing relatively high levels of HP and OP for activities such as video games, knowledge acquisition, and creative arts. Structural equation modeling showed that, as predicted by the DMP, HP was associated with positive emotions and flow. In turn, OP was positively associated with conflict, rumination, and negative emotions. Finally, HP and OP were positively and negatively associated with optimal functioning respectively.

**Conclusion:** Findings suggest that intense interests can be defined as passion using the DMP. The DMP offers a theoretical framework which can account for the duality of intense interest and predict psychological and functional outcomes. Learning to foster higher levels of HP for intense interests can improve well-being and promote positive psychological experiences.

**Keywords:** autism, passion, intense interests, well-being, optimal functioning

## Community brief

“Why is this an important issue?”

Autistic individuals have several passions which shape how they spend their time and what activities they engage in. Research and interventions in autism have historically focused on problematic aspects of passionate interests, often omitting the positive and motivational aspects of passion. Understanding the role that passion plays in their lives and how it relates to their well-being can inform clinical and research practices.

“What was the purpose of this study?”

Our goal was to study the intensity and the types of passion that autistic individuals showed for their passionate interests and how it related to their well-being. Using a well-established passion theory brought a new understanding to the field of autism research and passionate interests.

“What did the researchers do?”

We asked autistic teenagers and autistic young adults to fill out an online questionnaire about their current and most important passionate interest. They were asked to rate how passionate they felt about their interest, what emotions were associated with this interest, and how much they thought that passion conflicted with other aspects of their life. They also rated their global well-being and the quality of their relationships.

“What were the results of the study?”

We found that most autistic teenagers and young adults were highly passionate about their interest. We found evidence for high levels of both types of passion depending on the individual (obsessive and harmonious passion). Having more obsessive passion was associated with having lower well-being, and similarly more harmonious passion was related to greater well-being.

“What do these findings add to what was already known?”

We now know that passionate interests in autism can be defined as passions according to the dualistic model for passion theory. Both positive and negative impacts of passionate interests can coexist in one person. Passionate interests in autism are an important part of well-being and the type of passion is related either greater or lower well-being.

“What are potential weaknesses in the study?”

The study used multiple choice questionnaires designed for the general population, and which may not fully capture the lived experiences of autistic participants. Also, our study does not represent the entire spectrum of autism, as we excluded participants without reading and writing language skills.

“How will these findings help autistic adults now or in the future?”

Autistic individuals can evaluate their own passionate interests to better understand how different aspects of their passion are connected to their well-being. Also, we hope our results can help redefine passionate interests and bring empowerment, documenting the positive and motivational aspects of interests in autism.

**Association between passion and optimal functioning in autistic individuals: The Dualistic Model for Passion**

“It is a real obsession, which paradoxically, is also essential to maintain a good mood and to control my anxious thoughts associated with leading various aspects of my adult life.”

*21-year-old autistic woman who has a passion for education.*

The time and energy that autistic individuals invest in their passionate interests is substantial, holding its similarities to a passionate athlete’s rigorous workout schedule. In contrast to athletes, autistic individuals are not always valued for their interests, and some are discouraged from engaging in them through behavioral interventions.<sup>1-4</sup> Interventions aimed at diminishing or changing interests in autistic children are often motivated by the disruptions caused by the interest in everyday life and family functioning.<sup>2,5-7</sup> While autistic interests may come with insistence on sameness or adhering to routines, they have also been associated with positive emotions, intrinsic motivation, and well-being for autistic individuals over different age groups.<sup>7-9</sup> Interventions targeting interests have therefore been criticized<sup>10</sup>, namely for having failed to consider the positive aspects of interests and their functions for autistic individuals such as self-regulation and emotions.<sup>11,12</sup>

Intense interests, or restricted interests in the Diagnostic and Statistical Manual (DSM-5), are present in a majority of autistic individuals.<sup>13-15</sup> These interests can span from subjects such as music, history, and algebra, to video games. Some interests are known for being similar to those in the neurotypical population, with greater intensity.<sup>8,16</sup> Intense interests are reported as being more frequent in boys compared to girls<sup>17-19</sup> and are typically attributed to autistic

individuals without intellectual disability or language barriers.<sup>15</sup> They develop as early as three years of age and intensify throughout adulthood, along with other interests.<sup>14,20-22</sup>

Most of the existing measures assess negative aspects of interests (e.g.: disruption, frequency, and intensity of observed behaviors).<sup>26</sup> For example, diagnostic tools for autism such as the Autism Diagnostic Observation Schedule<sup>23</sup> (ADOS-2) and the Autism Diagnostic Interview<sup>24</sup> (ADI-R) rate the intensity and the disruption caused by these interests. While more measures now assess strengths and interests in children<sup>25-26</sup> and adults<sup>9</sup>, research contributing to an integrated conceptual understanding of motivational processes involved in interests and their impact on functioning is needed.

#### *Interests: Positive and Negative Aspects*

Positive impacts of intense interests range from social inclusion to learning, and positive emotions in autistic children (parent report), and in autistic adults (self-report).<sup>27-35</sup> Interests also serve as a coping strategy for autistic individuals when they experience negative emotions (e.g., in stressful situations) and can provide meaning to their environment.<sup>11,36</sup> A qualitative study interviewing autistic adults between the ages of 35 and 55 reported that some participants mentioned feeling at ease and confident in conversations when their area of expertise and interest was concerned.<sup>37</sup> Other studies also report that intense interests are part of autistic individual's identity and contribute to building self-confidence.<sup>33,38</sup> Finding peers that have similar interests is known as an easier way of making connections with others and developing friendships.<sup>35,39,40</sup>

According to research with autistic children and their parents, interests however can impede various aspects of daily functioning and social skill development.<sup>7,22,27,32,41</sup> Autistic children's interests were shown to be more intense and disruptive of family activities compared to non-autistic children.<sup>5-7</sup> During adolescence, intense interests have been shown to interfere

with personal short and long term goals, for example limiting choices having to do with a future career orientation.<sup>44</sup> Further, some authors proposed that intense interests may explain the lack of social interactions and interest toward other activities.<sup>7,43,45</sup> In Collis et al.<sup>12</sup>, autistic adults were asked to describe their interest and one of the emerging themes were negative aspects relating to social inclusion, namely because of stigma and others' perception. Some participants also reported having difficulties balancing their intense interest with other activities in their personal life, such as personal hygiene, sleep or social demands.<sup>12</sup>

### *Passion and Optimal Functioning*

The Dualistic Model for Passion (DMP) is a theoretical model that has led to a greater understanding of adaptive aspects of passion in the general population and specific groups of individuals (e.g.: athletes, nurses, students).<sup>46,47</sup> As there is evidence for both beneficial and detrimental characteristics of intense interests in autism, the DMP framework could contribute to understanding and predicting the impacts of interests on an individual level. According to the DMP, type and level of passion endorsed for activities or topics will predict positive and negative impacts on well-being and more largely, optimal functioning.<sup>46</sup>

Passion is a strong interest for an activity or topic that becomes part of one's identity and towards which substantial time and energy is invested.<sup>46,48</sup> Two types of passion are described by the DMP: harmonious passion (HP) and obsessive passion (OP). Individuals will express both types of passion at varying levels for one or more activities and topics. These two passions exist on different continua. HP is a flexible and willful engagement in an activity, free from external pressures (e.g., an activity practiced for its inherent benefits). HP is associated to intrinsic motivation and positive psychological experiences. OP is described as a strong need or an urge to engage in an activity, often in a rigid and compulsive manner.<sup>46,48</sup> It is associated with

environmental control (e.g.: practicing an activity for feelings of social acceptance or self-worth). Although OP may be associated with a greater sense of competence and expertise in an activity, it can also lead to lower well-being by fostering negative psychological outcomes (i.e., negative emotions, conflict).<sup>46,49,50</sup> Variable levels of HP and OP are found across individuals, and according to findings in the general population, having high levels of HP can be protective against the negative impacts of OP.<sup>51</sup>

Emotions, flow, rumination, and conflict are psychological processes that unfold when a person is pursuing a passion.<sup>46</sup> Specifically, positive emotions such as enthusiasm and pride are associated with a passion which has high levels of HP, whereas negative emotions such as irritability and anxiety are typically associated with OP.<sup>50,52,53</sup> Experiencing a state of flow (i.e., being completely immersed in the activity and having a sense of control),<sup>54</sup> is associated with HP.<sup>55</sup> In contrast, experiencing conflict and rumination is associated with OP.<sup>50,56</sup> Conflict arises when someone's passion interferes with important life engagements such as work, school, or relationships on a regular basis.<sup>56</sup> Rumination is the presence of thoughts related to the passionate activity outside of the activity practice.<sup>57</sup> These are non-intentional and recurrent thoughts that represent a person's preoccupations and unattained goals.<sup>57-58</sup> The above-mentioned psychological processes have an impact on optimal functioning, a multidimensional measure of happiness and functioning in the general population.<sup>59-60</sup> Specifically, optimal functioning comprises five dimensions: subjective well-being, physical health, quality of interpersonal relationships, performance in school or work, and contribution and implication to society via work, school or other means.<sup>49</sup> HP, positive emotions and flow positively predict optimal functioning, whereas OP, negative emotions, conflict, and rumination negatively predict optimal functioning.<sup>46,59-62</sup>

### *Bridging the Gap*

Previous studies in autism have established a connection between intense interests, flow, and well-being.<sup>8,63-66</sup> However, further exploration is needed to understand flow in the context of intense interests, as well as how it might be maintaining well-being. Measuring HP and OP and other variables within the DMP can serve to capture the duality of autistic adults' lived experiences of intense interests, without neglecting the positive aspects of flow and emotions. As mentioned by Champan and Carel<sup>67</sup> autistic individuals are oftentimes victims of systemic discrimination and are seen at odds with conceptions of well-being or thriving. The concept of optimal functioning is based on societal expectations and might not capture aspects of autistic flourishing or thriving concerning intense interests. Nonetheless, optimal functioning steps away from measures of "severity", "frequency", "disruption" and is interested in subjective well-being and other indicators of functioning in society (relational and physical health, performance, contribution to community). Understanding the role of autistic interests in emerging adults' optimal functioning remains relevant in the actual context, as autistic adults are at greater risk for discontinuing an academic program, have greater difficulty finding/maintaining stable employment and reaching parental independence, as well as experience lower social participation and quality of life.<sup>68-73</sup>

### *The Present Study*

The research objectives for this study were twofold. The first goal was to characterize passion in a population of autistic participants. Given that autistic individuals are known to have intense interests, they will likely be passionate about an activity or a subject, although it is yet to be determined whether their interests are more harmonious or obsessive. We expected passion levels to be high and hypothesized that high levels of obsessive passion would be present given the diagnostic criteria and measurement tools which insist on disruptiveness and rigid routines.<sup>13</sup> Concurrently, we expected levels of harmonious passion to be high given the flow and intrinsic motivation associated with intense interests in previous studies.<sup>8,35</sup>



The second aim was to determine whether HP and OP could predict emotions, flow, conflict, rumination, and optimal functioning. Based on the existing literature in non-autistic samples, we expected HP would be positively, whereas OP would be negatively associated, with optimal functioning (well-being, physical health, quality of interpersonal relationships, performance, and contribution to society). Similarly, HP was expected to be positively associated with positive emotions and flow, and negatively associated with negative emotions. OP was expected to be positively associated with negative emotions, conflict, and rumination.

## Materials and Methods

### *Participants and Procedure*

The present study is a cross-sectional online study with a final sample of 108 autistic individuals (43 men, 57 women, 7 non-binary, 1 missing), with a mean age of 23.66 years (SD=5.11 years; 14-33). Because participants' diagnosis could not be verified, researchers recruited via organizations working directly with autistic people in French speaking communities in Canada and France (e.g., Fédération québécoise de l'autisme). We provided a link in the advertisement and redirected participants to an online consent form and survey. Study materials were in French, and participants completed the survey without financial compensation. All surveys were completed anonymously on an online platform provided by the Université du Québec à Trois-Rivières (Chopin). Questionnaires were presented in the same sequence for all participants and total duration of the study was approximately 30 minutes. Participants were asked to report various information concerning their autism diagnosis (professional, age of diagnosis, etc.). We excluded participants reporting a self-diagnosis of autism ( $n = 2$ ). Co-occurring conditions were documented but did not represent an exclusion criterion. This study was approved by the Université du Québec à Trois-Rivières ethics board (CER-17-237-07.14).

Participants reported having been diagnosed by a medical doctor (63; 58.33%), or a psychologist (45; 41.67%), at a mean age of 17.60 years (SD=8.31). Participants' diagnoses were autism spectrum disorder (ASD; 26; 24%), Asperger's syndrome (50; 46.3%), Pervasive developmental disorder (5; 4.6%), autism (2; 1.9%) or ASD and Asperger's (25; 23.2%). Reported co-occurring conditions were common, with only 22% of the sample not reporting any other condition than autism ( $n=24$ ). In the current sample, 28.4% of participants reported having only one condition other than autism ( $n=31$ ), 21.1% reported having two ( $n=23$ ), and 28.5%

reported having from 3 to 5 co-occurring conditions (n=31). Of these, anxiety disorders, ADHD, and depression were the most reported, with 46, 35 and 22 participants respectively (see a detailed list of co-occurring conditions in Supplementary table 1). For additional demographic information see Table 1.

[insert table 1]

### *Measurements*

A sociodemographic questionnaire assessed age, gender, occupation, and other information related to the participants' autism diagnosis. Ethnicity was not measured to limit the duration of our online study. Unless otherwise specified, all variables were then measured using a 7-point Likert-type scale (scores ranging from 1 to 7). Higher scores indicated higher levels of the construct. Reliability coefficients were measured with Cronbach's alpha.

**Passion for an Activity.** Passion was measured using the Passion Scale, in which participants were asked to name and describe an activity that they loved, that they found important, and in which they invested time and energy (i.e., their passion).<sup>48,74</sup> This scale measures harmonious passion with six items (e.g., "My activity is well integrated in my life") and obsessive passion with six items (e.g., "I have difficulties controlling my urge to do my activity"). The Passion Scale has been used in the general population and was used for the first time with a sample of autistic individuals in this study. Satisfactory reliability was found for the HP ( $\alpha=.74$ ) and OP ( $\alpha=.80$ ). Five additional items serve to determine the presence of passion (i.e., passion criteria, e.g., I spend a lot of time doing this activity) ( $\alpha=.65$ ).

**Positive and Negative Affect.** The Positive and Negative Affect Schedule (PANAS) was used to measure positive and negative emotions during the activity.<sup>75</sup> Five positive emotions (interested, determined, alert, enthusiastic, active) and five negative emotions (anxious, upset,

hostile, irritable, nervous) were measured. Participants were asked to indicate how much they feel each of these emotions while doing their activity. This scale has been used in prior studies with samples of autistic individuals (alphas between .64 and .93).<sup>76,77</sup> Reliability in the current sample was adequate for negative emotions ( $\alpha=.78$ ) but not satisfactory for positive emotions ( $\alpha=.49$ ). Removing emotions “alert” and “active” increased reliability for positive emotions, although reliability for this scale was still poor ( $\alpha=.58$ ).

Flow. The Jackson and Marsh<sup>78</sup> flow scale was adapted to measure the flow during the activity and was used for the first time in an autistic population. Twelve items were selected for the study. An example item is: “When doing my favorite activity...I feel in total control of what I am doing”. The scale showed satisfactory reliability ( $\alpha = .86$ ).

Conflict. Developed by Vallerand et al.<sup>48</sup>, this scale assesses the level of conflict between the favorite activity and other daily life activities. It comprises five items and shows excellent psychometric properties in the general population.<sup>46,48,72</sup> A sample item is: “My activity conflicts with the other activities in my life”. The scale showed satisfactory reliability in the current sample ( $\alpha = .85$ ”).

Rumination. A 6-item scale was used to measure rumination outside of the favorite activity. Thinking about this activity while doing something else was measured with items such as: “When doing something outside of my favorite activity...I am constantly analyzing and thinking of my activity”. This scale was adapted from the Rumination on Sadness Scale<sup>79</sup>, was used in previous studies in the general population<sup>54</sup>, and showed excellent reliability in the current sample ( $\alpha=.90$ ).

Optimal Functioning in Society. Optimal functioning was assessed using the OFIS Scale.<sup>60</sup> With 35 items, the scale measures five different dimensions: psychological well-being,

physical health, performance at work or school, contribution to society, and interpersonal relationships. Four types of interpersonal relationships were measured and combined to form an average composite score: family, friendship, romantic, and an activity partner, if the participant's favorite activity was practiced with another person. For each relationship type, four items with a five-point Likert-type scale were used (0=not at all; 4=extremely) to evaluate how satisfying, harmonious, rewarding, and trustworthy the relationship was. Validation studies in the general population support OFIS scale's factorial structure (McDonald's Omega between .82 and .93 for each dimension) and show moderate to high item correlation for each dimension.<sup>49,59</sup> All five subscales showed good to excellent reliability in the current sample (performance,  $\alpha = .87$ ; well-being  $\alpha = .93$ ; physical health  $\alpha = .86$ ; contribution  $\alpha = .97$ ; interpersonal relationships,  $\alpha = .94$ ). Reliability for the overall optimal functioning construct was excellent in our sample ( $\alpha = .95$ ).

### *Statistical Analyses*

All analyses were performed in R version 4.0.5<sup>80</sup> at a significance criterion of  $p < .05$ . Preliminary analyses were conducted to examine outliers, missing data, and statistical assumptions (i.e., scale reliability, homoscedasticity, and univariate and multivariate normality). Descriptive analyses for aim 1 consisted of means, standard deviations, and Pearson correlations (pairwise). To answer the second aim, a path analysis model was performed using the *Lavaan* package.<sup>81</sup> Specifically, paths were specified from both HP and OP to optimal functioning, conflict, rumination, flow, positive and negative emotions. Full information maximum likelihood (FIML) was used in the model to account for patterns of missing data. The  $\chi^2$  statistic, the robust Comparative Fit Index (CFI), robust Tucker-Lewis Index (TLI), robust Root Mean Square Error of Approximation (RMSEA), Standardized Root Mean Square Residual (SRMR), Goodness-of-Fit Index (GFI) and Adjusted GFI were used to evaluate model fit. A statistically non-significant

$\chi^2$  statistic indicates that the model has a good fit. Furthermore, a CFI/TLI greater than .90 indicates an acceptable fit and a CFI/TLI greater than .95 indicates a good fit. The RMSEA was expected to be smaller or equal to 0.6 for a good model fit. Finally, a SRMR smaller than .10 indicates a good fit. GFI > .9 indicates a good fit, and the AGFI, adjusted for model parsimony, is expected to be greater than .5.

## Results

### *Preliminary Analyses*

There were missing data for a total of 20 participants. Four variables had more than 5% missing data (positive emotions = 8%; negative emotions = 8%; rumination = 9%; optimal functioning = 15%). Visual inspection of missing data suggested that this was due to attrition, as variables towards the end of the survey had more missing data. Further analyses on missing data indicated that they were missing completely at random (MCAR; Hawkins test  $p = .048$ ; non-parametric test  $p = .110$ ).<sup>81</sup> Univariate skewness and kurtosis indices were within the  $\pm 2$  margins, and all but two variables had indices within  $\pm 1$ , indicating univariate normality.<sup>83</sup> Multivariate normality was not respected, as indicated by the Mardia skewness and kurtosis tests ( $p < .05$ ). Robust estimator MLR was used for all path analyses because of the absence of multivariate normality in the sample. Variance Inflation Factor (VIF) was satisfactory, indicating no multicollinearity issues ( $< 2.50$ ).<sup>84,85</sup>

### *Descriptive Analyses*

Participants were passionate about various activities (see Table 2). They reported spending on average 25.34 hours per week ( $SD=16.67$ ) doing their favorite activity and have been practicing it for the last 10.58 years ( $SD=6.63$ ). Nearly half participants reported doing their activity alone ( $n=53$ ; 49.1%), or occasionally sharing this activity with another person ( $n=52$ ; 48.1%), and a minority always practiced their activity with other people ( $n=3$ ; 2.8%).

Two independent researchers categorized activities based on previous work from Vallerand et al. (study 1)<sup>48</sup>, showing satisfactory interrater reliability ( $\kappa = .83$ ). Categories were adapted to the current sample of participants, as the types of interests varied significantly from Vallerand and colleagues'<sup>48</sup> passion categories (ex.: gaming). Eight categories were


created: active arts ( $n = 22$ ), knowledge acquisition ( $n = 26$ ), gaming ( $n = 26$ ), television ( $n = 11$ ), reading ( $n = 11$ ), sports ( $n = 6$ ), animals ( $n = 4$ ), and passive arts ( $n = 2$ ). The most common categories in the present sample were gaming, knowledge acquisition and active arts (see Table 2). The most common interest for men was gaming ( $n = 20$ ), whereas women more commonly endorsed knowledge acquisition ( $n = 16$ ) and active arts ( $n = 14$ ). The category knowledge acquisition was defined as having a passion for a specific topic and accumulating facts and reading/researching on this topic. Within the category of knowledge acquisition, topics were classified under social sciences ( $n = 19$ , 10 men; sociology, philosophy, psychology, linguistics, history, education, politics) and sciences ( $n = 7$ , 1 man; medicine, biology, astronomy, neurology, physics, mathematics) in line with previous work in the field.<sup>7,86-89</sup>

[insert table 2]

All participants were highly passionate for their activity according to the passion criteria subscale (scores  $>4/7$ ; mean = 6.31), which is a greater prevalence of passion than in the general population, characterized by 15-30% of non-passionate individuals.<sup>48,61</sup> Harmonious and obsessive subscales revealed higher levels of HP (5.41/7) compared to OP (4.31/7; see Figure 1). Levels of HP are comparable to those found in previous studies, whereas OP levels are higher compared to the general population (see Supplementary table 2). Means and standard deviations for study variables are reported by gender in Supplementary table 3, and items means and standard deviations for each measure are available in supplementary tables 4-9.



Figure 1. Characterizing passion in autism




Means, standard deviations and correlations between variables are reported in Table 3. As expected, HP was positively and moderately associated with optimal functioning ( $r=.35$ ,  $p<.001$ ), whereas OP was strongly negatively associated with optimal functioning ( $r=-.44$ ,  $p<.001$ ). Moreover, HP was associated with experiencing flow ( $r=.26$ ,  $p=.008$ ) and positive emotions ( $r=.32$ ,  $p=.001$ ) during the activity, but also negatively associated with experiencing negative emotions ( $r=-.35$ ,  $p<.001$ ). Otherwise, OP was associated with experiencing greater negative emotions ( $r=.21$ ,  $p=.041$ ) during the activity, conflict ( $r=.79$ ,  $p<.001$ ) and rumination ( $r=.70$ ,  $p<.001$ ) outside of the activity. None of the variables were significantly related to the number of hours spent on the activity per week.

[insert table 3]

*Main Analyses*

With 40 parameters and 28 degrees of freedom, the model was under-identified. Paths were specified from HP and OP to optimal functioning, conflict, rumination, flow, positive and negative emotions (Figure 2). It was expected that HP would be positively, whereas OP would be negatively associated with optimal functioning. Similarly, HP was expected to be positively associated with positive emotions and flow, and negatively associated with negative emotions. OP was expected to be positively associated with negative emotions, conflict, and rumination. The model supported the hypotheses and yielded an acceptable fit to the data ( $\chi^2$  test:  $p = .291$ ; CFI = .995; TLI = .962; RMSEA = .055; SRMR = .048; GFI = .999; AGFI .992). Specifically, HP was positively associated with positive emotions ( $\beta = .347, p = .001$ ), flow ( $\beta = .250, p = .005$ ), and optimal functioning ( $\beta = .336$ ), and negatively associated with negative emotions ( $\beta = -.390, p = .001$ ). OP was positively linked to conflict ( $\beta = .780, p < .001$ ), rumination ( $\beta = .702, p < .001$ ), negative emotions ( $\beta = .158, p = .051$ ) but negatively associated with optimal functioning ( $\beta = -.430, p < .001$ ). Overall, the present results support the DMP in a sample of autistic participants.<sup>46</sup>

Figure 2. Structural equation modeling results



## Discussion

The present study investigated passion and optimal functioning in autism for the first time using the Dualistic Model for Passion (DMP).<sup>46,48</sup> The first research objective was to characterize intense interests by measuring harmonious passion (HP) and obsessive passion (OP) in autistic teenagers and young adults between ages 14 to 33. The second objective was to determine whether the two types of passion (i.e., HP and OP) were related to positive and negative emotions, flow, conflict, rumination and, optimal functioning (well-being, physical health, quality of interpersonal relationships, performance, and contribution to society). Based on the existing literature in non-autistic samples, it was expected that the type of passion would predict levels of optimal functioning, such that HP would be positively, whereas OP would be negatively associated, with optimal functioning. Similarly, HP was expected to be positively associated with positive emotions and flow, and OP was expected to be positively associated with negative emotions, conflict, and rumination.

Results indicated that autistic individuals are highly passionate for their interest, showing both high levels of harmonious and obsessive passion. As posited by the DMP, *the type* (i.e., HP and OP) *and intensity* of passion were predictive of optimal functioning and of the cognitive and emotional experience of intense interests in autism. HP positively predicted optimal functioning, flow, and positive emotions, while negatively predicting negative emotions. OP negatively predicted optimal functioning, while positively predicting conflict and rumination. These results are discussed in the light of current findings in the passion and autism literatures.

### *Describing Passions in Autism*

All participants in the sample were passionate for their interest, with HP being more present than OP (means of 5.41 vs 4.31, respectively). Similarly, studies measuring passion using

the Passion Scale<sup>48,74</sup> in the general population find higher levels of HP compared to OP.<sup>61,89</sup> HP levels found in the current sample are similar to those found in the general population (5.26-5.92).<sup>58,85</sup> In contrast, OP levels are typically lower in the general population (3.10-3.69).<sup>89,90</sup> To note, OP levels found in the present study resemble those of elite athletes<sup>91</sup> (i.e., 4.78 for OP), which are higher than most samples from the general population. Moreover, autistic individuals were found to spend more time on their passion (25.5 h/week) compared to the general population, typically spending on average from 8 to 15.5 h/week on their interest.<sup>48,61,74,92</sup> As such, passions in autism as harmonious as passions in the general population and can be described as more obsessive and intense in terms of time investment. Accordingly, past research with autistic participants has found positive experiences are tied to passions but these experiences can come at a cost for general well-being in certain situations or individuals. Specifically, in Rapaport et al.<sup>64</sup>, semi-structured interviews with 24 autistic adults indicated that although practicing the interest is pleasant, exhaustion or losing oneself in the activity was a frequent experience. Losing control over one's activity and having difficulty disengaging from the task is compatible with descriptions of OP. Both types of passions help understand the different lived experiences of autistic adults and their intense interests, as one can simultaneously experience the benefits of harmonious passion but endure the consequences of obsessive passion. Impacts on well-being and optimal functioning will therefore depend on the level of both types of passion within one individual at a given time.

Previous research on interests in autism have described an "autistic flow" and mentioned that positive emotions were commonly experienced.<sup>9,38</sup> Adding to the literature, current findings suggest that within an autistic sample, flow and emotions will vary according to the type of passion which is most endorsed. Specifically, flow states and positive emotions are associated

with the presence of HP and not merely the presence of passion, as OP was not found to correlate with flow states nor with positive emotions. As HP is characterized by a non-defensive form of engagement which comes with openness to experience, a secure self-esteem and mindfulness, it can lead to a state of flow more readily than OP.<sup>46</sup> High levels of OP lead to lower levels of flow and more negative emotions because the passion is internally controlled. It leads to a person having an eye on the task but the other on external elements or self-evaluation (e.g.: the expected outcomes).<sup>46</sup> What remains uncertain is whether OP can account for certain aspects of intense interests which are reported in the literature (e.g., having trouble disengaging from a pleasant task).<sup>64</sup> Namely, the autistic tendency to selectively focus on one thing at a time (i.e., monotropism) can induce hyperfocus states or autistic flow states, which differ from the classic definition of flow and could correspond with OP.<sup>65,66</sup> For example, having difficulty disengaging from a task is not associated with flow but more likely attributable to other cognitive (e.g., monotropism, executive functioning) or possibly motivational factors (e.g., OP). Typically, flow states are not compatible with descriptions of individuals losing track of time to a point where basic needs (e.g., eating, sleeping) conflict with the interest.<sup>54</sup>

The current study suggests that HP could account for some of the adaptive consequences of intense interests, whereas OP would account for less adaptive outcomes. Namely, previous findings in autism found that conflict and interference with daily activities or family activities were characteristic of intense interests.<sup>5,6,12,14</sup> While conflict or interference may vary with topics for example, it may also vary according to OP levels. The current study found negative impacts related to intense interest but also considered individual differences in passion levels. Higher levels of OP were positively associated with negative emotions, conflict, and rumination. According to the DMP, OP is characterized by rigidity and persistence and will lead to conflict

with other areas of a person's life.<sup>46</sup> It is also accompanied by a contingent self-esteem, which entails that a person's self-esteem and emotions will fluctuate according to how well their passion is progressing (e.g.: winning a game, reaching a target). As a result, individuals with high OP tend to ruminate more on their passion compared to those with high HP.

A diversity of topics was found in the sample, with video games, active arts (ex.: painting, singing, acting) and knowledge acquisition (learning things about a specific topic) being the most common themes. These categories differ from the ones identified in the general population, which were commonly sports and music.<sup>48</sup> Potentially, as described in the Monotropism theory<sup>66</sup>, autistic people would be more easily drawn towards less social endeavors, where things are more predictable. Additionally, having an area of expertise provides a sense of control and can help feel secure when talking about the topic of expertise to other people.<sup>12</sup> Interests are commonly used by autistic individuals to form social bonds.<sup>63</sup> In Turner-Brown et al.<sup>7</sup>, autistic children also strived for peer acceptance by showing off their knowledge and hoped their expertise would be recognized by others.

Women in the sample were more drawn to knowledge acquisition and active arts, and men reported their main interest as being video games more than any other activity. The category of knowledge acquisition was subcategorized into topics of the social sciences (ex.: history, politics, linguistics) and the sciences (ex.: physics, mathematics, biology), following previous work in the field<sup>86,87</sup>. Social sciences were more commonly endorsed in the present sample, although gender was equally distributed across social sciences and sciences topics. This replicates more recent work in children with autism, in which autistic girls and boys did not show a gender difference.<sup>87</sup>

*Does Passion Lead to Optimal Functioning?*

The current study showed that autistic individuals' passions were related to the five dimensions of optimal functioning (i.e., psychological well-being, physical health, relationship quality, contribution to society, performance at work or school). In agreement to Vallerand's perspective, the quantity and quality of passion (HP or OP) predicted psychological well-being in autistic individuals.<sup>46</sup> Engaging in pleasant activities is not sufficient to encourage optimal functioning, rather people need to pursue their interests out of HP, rather than OP.<sup>47</sup> Higher levels of HP were associated with greater optimal functioning, whereas higher levels of OP were related to lower optimal functioning in autistic young adults and teenagers. In contrast with Grove et al's<sup>8</sup> findings in the autism literature, the time invested in the activity itself (i.e., number of hours per week) was not associated with well-being or other components of optimal functioning. Therefore, the present results suggest that finding a balance between HP and OP may have a greater influence on well-being compared to the amount of time spent on the interest.

Even though OP was associated with lower optimal functioning than HP, it nonetheless has been found to positively relate to basic psychological needs.<sup>92</sup> Because of this, even reducing a highly obsessive interest would reduce a person's satisfaction in life and would potentially decrease their optimal functioning. Holding et al.<sup>93</sup> hypothesized that behavioral addiction could develop when a person's needs in the rest of one's life (outside one's major interest) are not satisfied. OP would represent a compensatory mechanism for the person to fulfill one's needs within the passionate activity.<sup>93</sup> Coutelle et al.<sup>94</sup> argues that intense interests in autistic individuals "should be preserved", even in cases of video gaming, where the line between addiction and interest is unclear. Coutelle et al. 's<sup>94</sup> recommendations include limiting and monitoring usage of video games in cases of addiction and investing time in other activities



outside of gaming. Thus, the addiction and passion literature recommend finding new activities. Consequently, the passionate activity would seem like less of a necessity (i.e., less obsessive).

Passionate people generally have greater optimal functioning, particularly those with high HP.<sup>51,61</sup> Paradoxically, although autistic adults were found to be highly passionate, they generally have lower well-being and quality of life compared to non-autistics.<sup>95</sup> This well-being gap between autistic and non-autistic adults has been shown to occur in varying degrees depending on factors such as intellectual functioning, presence of other diagnoses, or experiencing social stigma.<sup>67,93</sup> Specific aspects of everyday life need to be considered when measuring well-being or quality of life in autistic adults (e.g., sensory issues, stigmatization)<sup>96,97</sup> and available measures may not capture aspects of autistic thriving that are not valued or attended to in the general population.<sup>67</sup> Having a harmonious or obsessive passion could be one of those contributing aspects, as harmonious passion contributes both to hedonistic well-being (i.e., having positive emotions) and eudaimonic well-being (i.e., finding meaning in life).<sup>61</sup>

### *Limitations*

This study has several limitations. First, this study was conducted online which limits the scope of generalizability of results. Second, a disproportionate amount of women are present in the sample as compared to clinical samples of autism which refer to 1:4 girls:boys.<sup>98</sup> Third, self-report questionnaires were used, and they limit the participation for individuals with intellectual disabilities or communication barriers. Furthermore, they are subjective in nature and should be corroborated in future research while using objective measures or through observation. Our study invited participants to select only one of their favorite activities, which can limit the representativeness of the findings, and which could also have biased participants towards choosing a favorite interest which was more harmonious. Additionally, ethnicity data was not

collected in the current study which can also limit the generalizability of the findings. Finally, the present study used a cross-sectional design, and it cannot lead to causality inferences.

#### *Future Directions*

Considering current limitations, it would be important to investigate passion using different designs and methodologies. First, interviews with autistic adults could be conducted to gain deeper insight into the nature of their passion and its consequences on optimal functioning. Second, a longitudinal design would help to understand the directionality of the relations found between passion, the different psychological processes associated with passion and optimal functioning. Lastly, positive psychology can offer relevant theoretical models for investigating the processes through which passion leads to well-being and quality of life in autism.<sup>99</sup>

#### **Conclusion**

Investigating passion in autistic individuals can contribute to a better understanding of the duality of psychological impacts associated with intense interests. Using the Dualistic Model for Passion (DMP) to study intense interests revealed that autistic teenagers and young adults' interests can be described by measuring levels of harmonious and obsessive passion. The most common interests were gaming, knowledge acquisition, and artistic pastimes. Importantly, the number of hours spent engaging in their passion was substantial (25 hours) but was not associated with either harmonious or obsessive passion. Harmonious passion was associated with positive emotions, flow, and optimal functioning, whereas obsessive passion was positively related to negative emotions, rumination, conflict, and lower levels of optimal functioning. The current study suggests that the duality of intense interests can be measured using the DMP, which can help predict impacts on well-being and optimal functioning in emerging autistic

adults. Future research on the role of passion in autism seems promising as to reverse the stigma associated with intense interests and promote positive perspectives.

### **Acknowledgements**

The authors thank research participants for their time and essential contribution to this research project. We would also like to thank Delphine Périard-Larivée for her assistance with data collection, Marie-Pier Vaillancourt-Morel and Hugues Leduc for their guidance with statistical analyses, and Audrée-Anne Dumas for the revision of the final draft.

### **Authorship confirmation/contribution statement**

A.J.M. ran the data collection, analyzed the data with N.C. and wrote the original manuscript. E.-L.B, I.S. and R.J.V took part in conceptualisation and designed the methodology of the study. N.C., R.J.V., M.C., E.G. contributed significantly to revisions of the article, and E.-L.B. and I.S. offered supervision of the research activities. All authors confirm their original contributions to the article and reviewed and approved the full article before submission.

### **Statements and Declarations**

The research leading to these results received funding from the Social Sciences and Humanities Research Council (SSHRC) under Grant Agreement No 430-2020-0669 and the Regroupement Intersectoriel de recherche en Santé de l'Université du Québec (RISUQ; Pilot project grant). The authors have no relevant financial or non-financial interests to disclose and no competing interests to declare.

### References

1. Leekam SR, Prior MR, Uljarevic M. Restricted and repetitive behaviors in autism spectrum disorders: a review of research in the last decade. *Psychological bulletin*. 2011;137(4):562; doi : 10.1037/a0023341
2. Lin CE, Koegel R. Treatment for Higher-Order Restricted Repetitive Behaviors (H-RRB) in Children with Autism Spectrum Disorder. *Journal of Autism and Developmental Disorders*. 2018;48(11):3831-3845; doi: 10.1007/s10803-018-3637-3
3. Mercier C, Mottron L, Belleville S. A psychosocial study on restricted interests in high functioning persons with pervasive developmental disorders. *Autism*. 2000;4(4):406-425; doi:10.1177/1362361300004004006
4. Mottron L. Should we change targets and methods of early intervention in autism, in favor of a strengths-based education? *European child & adolescent psychiatry*. 2017;26(7):815-825; doi: 10.1007/s00787-017-0955-5
5. Anthony LG, Kenworthy L, Yerys BE, et al. Interests in high-functioning autism are more intense, interfering, and idiosyncratic than those in neurotypical development. *Development and psychopathology*. 2013;25(3):643-652; doi:10.1017/S0954579413000072
6. Cascio CJ, Foss-Feig JH, Heacock J, et al. Affective neural response to restricted interests in autism spectrum disorders. *Journal of Child Psychology and Psychiatry*. 2014;55(2):162-171; doi.org:10.1111/jcpp.12147
7. Turner-Brown LM, Lam KS, Holtzclaw TN, Dichter GS, Bodfish JW. Phenomenology and measurement of circumscribed interests in autism spectrum disorders. *Autism*. 2011;15(4):437-456; doi.org/10.1177/1362361310386507

8. Grove R, Hoekstra RA, Wierda M, Begeer S. Special interests and subjective wellbeing in autistic adults. *Autism Research*. 2018;11(5):766-775; doi.org/10.1002/aur.1931
9. Grove R, Roth I, Hoekstra RA. The motivation for special interests in individuals with autism and controls: Development and validation of the special interest motivation scale. *Autism Research*. 2016;9(6):677-688; doi:10.1002/aur.1560
10. Schuck RK, Tagavi DM, Baiden KM, et al. Neurodiversity and Autism Intervention: Reconciling Perspectives Through a Naturalistic Developmental Behavioral Intervention Framework. *Journal of Autism and Developmental Disorders*. 2021:1-21; doi:10.1007/s10803-021-05316-x
11. Lung SLM, Picard È, Soulières I, Bertone A. Identifying the functions of restricted and repetitive behaviours and interests in Autism: A scoping review. *Research in Autism Spectrum Disorders*. 2024;117:102458.
12. Collis E, Gavin J, Russell A, Brosnan M. Autistic adults' experience of restricted repetitive behaviours. *Research in Autism Spectrum Disorders*. 2022;90:101895; doi: 10.1016/j.rasd.2021.101895
13. American Psychiatric Association D. *Diagnostic and statistical manual of mental disorders: DSM-5*. Washington, DC: American psychiatric association; 2013.
14. Nowell KP, Bernardin CJ, Brown C, Kanne S. Characterization of special interests in autism spectrum disorder: A brief review and pilot study using the special interests survey. *Journal of Autism and Developmental Disorders*. 2021;51(8):2711-2724; doi:10.1007/s10803-020-04743-6
15. Spackman E, Smillie LD, Frazier TW, et al. Characterizing restricted and unusual interests in autistic youth. *Autism Research*. 2023;16(2):394-405; doi:10.1002/aur.2863

16. Cho IY, Jelinkova K, Schuetze M, et al. Circumscribed interests in adolescents with Autism Spectrum Disorder: A look beyond trains, planes, and clocks. *PloS one*. 2017;12(11):e0187414. Doi:10.1371/journal.pone.0187414
17. Frazier TW, Georgiades S, Bishop SL, Hardan AY. Behavioral and cognitive characteristics of females and males with autism in the Simons Simplex Collection. *Journal of the American Academy of Child & Adolescent Psychiatry*. 2014;53(3):329-340. e323; doi :10.1016/j.jaac.2013.12.004
18. Hartley SL, Sikora DM. Sex differences in autism spectrum disorder: an examination of developmental functioning, autistic symptoms, and coexisting behavior problems in toddlers. *Journal of autism and developmental disorders*. 2009;39:1715-1722; doi:10.1007/s10803-009-0810-8
19. Szatmari P, Liu XQ, Goldberg J, et al. Sex differences in repetitive stereotyped behaviors in autism: implications for genetic liability. *American Journal of Medical Genetics Part B: Neuropsychiatric Genetics*. 2012;159(1):5-12; doi:10.1002/ajmg.b.31238
20. Seltzer MM, Krauss MW, Shattuck PT, Orsmond G, Swe A, Lord C. The symptoms of autism spectrum disorders in adolescence and adulthood. *Journal of autism and developmental disorders*. 2003;33(6):565-581; doi:10.1023/B:JADD.0000005995.02453.0b
21. Shattuck PT, Seltzer MM, Greenberg JS, et al. Change in autism symptoms and maladaptive behaviors in adolescents and adults with an autism spectrum disorder. *Journal of autism and developmental disorders*. 2007;37(9):1735-1747; doi:10.1007/s10803-006-0307-7

22. South M, Ozonoff S, McMahon WM. Repetitive behavior profiles in Asperger syndrome and high-functioning autism. *Journal of autism and developmental disorders*. 2005;35(2):145-158; doi:10.1007/s10803-004-1992-8
23. Lord C, Rutter M, DiLavore P, Risi S, Gotham K, Bishop S. Autism diagnostic observation schedule—2nd edition (ADOS-2). *Los Angeles, CA: Western Psychological Corporation*. 2012;284.
24. Rutter M, Le Couteur A, Lord C. Autism diagnostic interview-revised. *Los Angeles, CA: Western Psychological Services*. 2003;29(2003):30.
25. Smerbeck A. The Survey of Favorite Interests and Activities: Assessing and understanding restricted interests in children with autism spectrum disorder. *Autism*. 2019;23(1):247-259; doi:10.1177/1362361317742140
26. Larose V, Sotelo K, Mottron L, Jacques C. Initial development of a questionnaire about parents' perspectives on the strengths and interests of autistic preschoolers. *Canadian Journal of Behavioural Science/Revue canadienne des sciences du comportement*. 2021;53(4):530-535; doi:10.1037/cbs0000268
27. Attwood A. *The complete guide to Asperger's syndrome*. Jessica Kingsley Publishers; 2006.
28. Bennett E, Heaton P. Is talent in autism spectrum disorders associated with a specific cognitive and behavioural phenotype? *Journal of autism and developmental disorders*. 2012;42(12):2739-2753; doi:10.1007/s10803-012-1533-9
29. Boyd BA, Conroy MA, Mancil GR, Nakao T, Alter PJ. Effects of circumscribed interests on the social behaviors of children with autism spectrum disorders. *Journal of autism and developmental disorders*. 2007;37(8):1550-1561; doi:10.1007/s10803-006-0286-8

30. Davey L. Using the special interests of autistic children to facilitate meaningful engagement and learning. *Good Autism Practice (GAP)*. 2020;21(1):43-64.
31. Gunn KC, Delafield-Butt JT. Teaching children with autism spectrum disorder with restricted interests: A review of evidence for best practice. *Review of Educational Research*. 2016;86(2):408-430; doi:10.3102/0034654315604027
32. Klin A, Danovitch JH, Merz AB, Volkmar FR. Circumscribed interests in higher functioning individuals with autism spectrum disorders: An exploratory study. *Research and Practice for Persons with Severe Disabilities*. 2007;32(2):89-100; doi:10.2511/rpsd.32.2.89
33. Patten Koenig K, Hough Williams L. Characterization and utilization of preferred interests: A survey of adults on the autism spectrum. *Occupational Therapy in Mental Health*. 2017;33(2):129-140; doi:10.1080/0164212X.2016.1248877
34. Sasson NJ, Dichter GS, Bodfish JW. Affective responses by adults with autism are reduced to social images but elevated to images related to circumscribed interests. *PLoS ONE*. 2012; 7, e42457; doi:10.1371/journal.pone.0042457
35. Wood R. Autism, intense interests and support in school: from wasted efforts to shared understandings. *Educational Review*. 2021;73(1):34-54; doi:10.1080/00131911.2019.1566213
36. Spiker MA, Lin CE, Van Dyke M, Wood JJ. Restricted interests and anxiety in children with autism. *Autism*. 2012;16(3):306-320; doi :10.1177/1362361311401763
37. Manor-Binyamini I, Schreiber-Divon M. Repetitive behaviors: Listening to the voice of people with high-functioning autism spectrum disorder. *Research in Autism Spectrum Disorders*. 2019;64:23-30. Doi:10.1016/j.rasd.2019.04.001



38. Winter-Messiers MA. From tarantulas to toilet brushes: Understanding the special interest areas of children and youth with Asperger syndrome. *Remedial and Special Education*. 2007;28(3):140-152; doi:10.1177/07419325070280030301
39. Chan DV, Doran JD, Galobardi OD. Beyond friendship: The spectrum of social participation of autistic adults. *Journal of Autism and Developmental Disorders*. 2023;53(1):424-437; doi:10.1007/s10803-022-05441-1
40. Sosnowy C, Silverman C, Shattuck P, Garfield T. Setbacks and successes: How young adults on the autism spectrum seek friendship. *Autism in Adulthood*. 2019;1(1):44-51; doi:10.1089/aut.2018.0009
41. Ghanouni P, Jarus T, Zwicker JG, Lucyshyn J, Chauhan S, Moir C. Perceived barriers and existing challenges in participation of children with autism spectrum disorders: “He did not understand and no one else seemed to understand him”. *Journal of autism and developmental disorders*. 2019;49(8):3136-3145; doi:10.1007/s10803-019-04036-7
42. Lam KS, Bodfish JW, Piven J. Evidence for three subtypes of repetitive behavior in autism that differ in familiarity and association with other symptoms. *Journal of Child Psychology and Psychiatry*. 2008;49(11):1193-1200; doi:10.1111/j.1469-7610.2008.01944.x
43. Pierce K, Courchesne E. Evidence for a cerebellar role in reduced exploration and stereotyped behavior in autism. *Biological psychiatry*. 2001;49(8):655-664; doi:10.1016/S0006-3223(00)01008-8
44. Hodgetts S, Richards K, Park E. Preparing for the future: multi-stakeholder perspectives on autonomous goal setting for adolescents with autism spectrum disorders. *Disability and rehabilitation*. 2018;40(20):2372-2379; doi:10.1080/09638288.2017.1334836

45. Attwood A. *Asperger's syndrome: A guide for parents and professionals*. Jessica Kingsley Publishers; 1997.
46. Vallerand RJ. *The psychology of passion: A dualistic model*. Series in Positive Psychology; 2015.
47. Vallerand RJ, Robichaud J-M, Rahimi S, Bélanger JJ. Who's the Happiest and Why? The role of passion and self-regulation in psychological well-being. *Motivation and Emotion*. 2024;1-26; doi:10.1007/s11031-024-10069-y
48. Vallerand RJ, Blanchard C, Mageau GA, et al. Les passions de l'ame: on obsessive and harmonious passion. *Journal of personality and social psychology*. 2003;85(4):756; doi:10.1037/0022-3514.85.4.756
49. Bragoli-Barzan L, Vallerand R. Bonheur et engagement: le rôle de la passion dans le fonctionnement optimal en société. *Revue québécoise de psychologie*. 2017;38(2):21-41; doi :10.7202/1040769ar
50. Philippe FL, Vallerand RJ, Lavigne GL. Passion does make a difference in people's lives: A look at well-being in passionate and non-passionate individuals. *Applied Psychology: Health and Well-Being*. 2009;1(1):3-22; doi:10.1037/a0018017
51. Schellenberg BJ, Verner-Filion J, Gaudreau P, Bailis DS, Lafrenière MAK, Vallerand RJ. Testing the dualistic model of passion using a novel quadripartite approach: A look at physical and psychological well-being. *Journal of personality*. 2019;87(2):163-180; doi:10.1111/jopy.12378
52. Rousseau FL, Vallerand RJ. An examination of the relationship between passion and subjective well-being in older adults. *The International Journal of Aging and Human Development*. 2008;66(3):195-211; doi:10.2190/AG.66.3.b

53. Waugh CE, Fredrickson BL. Nice to know you: Positive emotions, self–other overlap, and complex understanding in the formation of a new relationship. *The journal of positive psychology*. 2006;1(2):93-106; doi:10.1080/17439760500510569
54. Csikszentmihalyi M. The flow experience and its significance for human psychology. *Optimal experience: Psychological studies of flow in consciousness*. 1988;2:15-35.
55. Carpentier J, Mageau GA, Vallerand RJ. Ruminations and flow: Why do people with a more harmonious passion experience higher well-being? *Journal of Happiness studies*. 2012;13(3):501-518; doi:10.1007/s10902-011-9276-4
56. Vallerand RJ, Paquet Y, Philippe FL, Charest J. On the role of passion for work in burnout: A process model. *Journal of personality*. 2010;78(1):289-312.
57. Martin, L. L., Tesser, A. Some ruminative thoughts. *Advances in Social Cognition*. 1996;9, 1–47.
58. Klinger, E. Consequences of commitment to and disengagement from incentives. *Psychological Review*. 1975;82(1), 1–25; doi:10.1037/h0076171
59. Bouizegarene N, Bourdeau S, Leduc C, Gousse-Lessard A-S, Houlfort N, Vallerand RJ. We are our passions: The role of identity processes in harmonious and obsessive passion and links to optimal functioning in society. *Self and Identity*. 2018;17(1):56-74; doi:10.1080/15298868.2017.1321038
60. Vallerand RJ. Passion and optimal functioning in society: a eudaimonic perspective. In A. S. Waterman (Ed.), *The best within us: Positive psychology perspectives on eudaimonia* (pp. 183–206). 2013. American Psychological Association. Doi:10.1037/14092-010

61. Philippe FL, Vallerand RJ, Lavigne GL. Passion does make a difference in people's lives: A look at well-being in passionate and non-passionate individuals. *Applied Psychology: Health and Well-Being*. 2009;1(1):3-22; doi:10.1111/j.1758-0854.2008.01003.x
62. Schellenberg BJ, Bailis DS. Can passion be polyamorous? The impact of having multiple passions on subjective well-being and momentary emotions. *Journal of Happiness Studies*. 2015;16(6):1365-1381; doi:10.1007/s10902-014-9564-x
63. Dwyer P, Williams ZJ, Lawson WB, Rivera SM. A trans-diagnostic investigation of attention, hyper-focus, and monotropism in autism, attention dysregulation hyperactivity development, and the general population. *Neurodiversity*. 2024;2; doi:27546330241237883.
64. Rapaport H, Clapham H, Adams J, Lawson W, Porayska-Pomsta K, Pellicano E. “In a State of Flow”: A Qualitative Examination of Autistic Adults' Phenomenological Experiences of Task Immersion. *Autism in Adulthood*. 2023; doi:10.1089/aut.2023.0032
65. Heasman B, Williams G, Charura D, Hamilton LG, Milton D, Murray F. Towards autistic flow theory: A non-pathologising conceptual approach. *Journal for the Theory of Social Behaviour*. 2024; 1-29; doi:10.1111/jtsb.12427
66. Lawson W. *The Passionate Mind: How People with Autism Learn*. Jessica Kingsley Publishers; 2011.
67. Chapman R, Carel H. Neurodiversity, epistemic injustice, and the good human life. *Journal of Social Philosophy*. 2022.
68. Ayres, M., Parr, J. R., Rodgers, J., Mason, D., Avery, L., & Flynn, D. A systematic review of quality of life of adults on the autism spectrum. *Autism*. 2018;22(7), 774–783; doi:10.1177/1362361317714988

69. Bishop-Fitzpatrick L, Minshew NJ, Mazefsky CA, Eack SM. Perception of life as stressful, not biological response to stress, is associated with greater social disability in adults with autism spectrum disorder. *Journal of autism and developmental disorders*. 2017;47:1-16; doi:10.1007/s10803-016-2910-6
70. Graham Holmes L, Zampella CJ, Clements C, et al. A lifespan approach to patient-reported outcomes and quality of life for people on the autism spectrum. *Autism Research*. 2020;13(6):970-987; doi:10.1002/aur.2275
71. Roux AM, Shattuck PT, Rast JE, et al. Characteristics of Two-Year College Students on the Autism Spectrum and Their Support Services Experiences. *Autism Research and Treatment*. 2015;2015(1):391693; doi:10.1155/2015/391693
72. Taylor EC, Livingston LA, Clutterbuck RA, Callan MJ, Shah P. Psychological strengths and well-being: Strengths use predicts quality of life, well-being and mental health in autism. *Autism*. 2023:13623613221146440; doi:10.1177/13623613221146440
73. Vincent A, Da Fonseca D, Baumstarck K, Charvin I, Alcaraz-Mor R, Lehucher-Michel M-P. The quality of life and the future of young adults with Asperger syndrome. *Disability and Rehabilitation*. 2020;42(14):1987-1994;doi:10.1080/09638288.2018.1544297
74. Marsh HW, Vallerand RJ, Lafrenière M-AK, et al. Passion: Does one scale fit all? Construct validity of two-factor passion scale and psychometric invariance over different activities and languages. *Psychological assessment*. 2013;25(3):796; doi :10.1037/a0032573

75. Watson D, Clark LA, Tellegen A. Development and validation of brief measures of positive and negative affect: the PANAS scales. *Journal of personality and social psychology*. 1988;54(6):1063; doi:10.1037/0022-3514.54.6.1063
76. Rueda P, Fernández-Berrocal P, Schonert-Reichl KA. Perspective-taking and empathic concern as mediators for happiness and positive affect in adolescents with and without asperger syndrome. *Journal of Developmental and Physical Disabilities*. 2014;26(6):717-735; doi:10.1007/s10882-014-9391-3
77. Samson AC, Huber O, Gross JJ. Emotion regulation in Asperger's syndrome and high-functioning autism. *Emotion*. 2012;12(4):659; doi:10.1037/a0027975
78. Jackson SA, Marsh HW. Development and validation of a scale to measure optimal experience: The Flow State Scale. *Journal of sport and exercise psychology*. 1996;18(1):17-35; doi:10.1123/jsep.18.1.17
79. Conway M, Csank PA, Holm SL, Blake CK. On assessing individual differences in rumination on sadness. *Journal of personality assessment*. 2000;75(3):404-425; doi:10.1207/S15327752JPA7503\_04
80. Team RC. R: a language and environment for statistical computing. Vienna: R Foundation for Statistical Computing. 2021.
81. Rosseel Y. lavaan: An R package for structural equation modeling. *Journal of statistical software*. 2012;48:1-36.
82. Jamshidian M, Jalal S. Tests of homoscedasticity, normality, and missing completely at random for incomplete multivariate data. *Psychometrika*. 2010;75(4):649-674; doi:10.1007/s11336-010-9175-3
83. Bulmer, M. G. Principles of statistics. Courier Corporation. 1979.

84. Myers RH, Myers RH. *Classical and modern regression with applications*. Vol 2: Duxbury press Belmont, CA; 1990.
85. Pituch KA, Stevens JP. *Applied multivariate statistics for the social sciences: Analyses with SAS and IBM's SPSS*. Routledge; 2015; doi:10.4324/9781315814919
86. Baron-Cohen S, Wheelwright S. 'Obsessions' in children with autism or Asperger syndrome: Content analysis in terms of core domains of cognition. *The British Journal of Psychiatry*. 1999;175(5):484-490.
87. Nowell SW, Jones DR, Harrop C. Circumscribed interests in autism: Are there sex differences? *Advances in Autism*. 2019;5(3):187-198; doi:10.1108/AIA-09-2018-0032
88. Caldwell-Harris CL, Jordan CJ. Systemizing and special interests: Characterizing the continuum from neurotypical to autism spectrum disorder. *Learning and Individual Differences*. 2014;29:98-105.
89. Stenseng F, Forest J, Curran T. Positive emotions in recreational sport activities: The role of passion and belongingness. *Journal of Happiness Studies*. 2015;16:1117-1129; doi:10.1007/s10902-014-9547-y
90. Philippe FL, Vallerand RJ, Andrianarisoa J, Brunel P. Passion in Referees: Examining Their Affective and Cognitive Experiences in Sport Situations. *Journal of Sport and Exercise Psychology*. 2009;31(1):77-96. doi:10.1123/jsep.31.1.77
91. Verner-Filion J, Vallerand RJ, Amiot CE, Mocanu I. The two roads from passion to sport performance and psychological well-being: The mediating role of need satisfaction, deliberate practice, and achievement goals. *Psychology of Sport and Exercise*. 2017;30:19-29; doi:10.1016/j.psychsport.2017.01.009

92. Lalande D, Vallerand RJ, Lafrenière MAK, et al. Obsessive passion: A compensatory response to unsatisfied needs. *Journal of personality*. 2017;85(2):163-178; doi:10.1111/jopy.12229
93. Holding AC, Verner-Filion J, Lalande D, Schellenberg BJ, Vallerand RJ. The roles of need satisfaction and passion in symptoms of behavioral addiction: The case of video gaming and gambling. *Motivation Science*. 2021;7(3):345; doi :10.1037/mot0000241
94. Coutelle R, Weiner L, Paasche C, et al. Autism Spectrum Disorder and Video Games: Restricted Interests or Addiction? *International Journal of Mental Health and Addiction*. 2021; doi:10.1007/s11469-021-00511-4
95. Lawson LP, Richdale AL, Haschek A, et al. Cross-sectional and longitudinal predictors of quality of life in autistic individuals from adolescence to adulthood: The role of mental health and sleep quality. *Autism*. 2020;24(4):954-967; doi:10.1177/1362361320908107
96. McConachie H, Mason D, Parr JR, Garland D, Wilson C, Rodgers J. Enhancing the Validity of a Quality of Life Measure for Autistic People. *Journal of Autism and Developmental Disorders*. 2018;48(5):1596-1611; doi:10.1007/s10803-017-3402-z
97. Caron V, Jeanneret N, Giroux M, et al. Sociocultural context and autistics' quality of life: A comparison between Québec and France. *Autism*. 2022;26(4):900-913; doi:10.1177/13623613211035229
98. Rødgaard EM, Jensen K, Miskowiak KW, Mottron L. Representativeness of autistic samples in studies recruiting through social media. *Autism Research*. 2022;15(8):1447-1456; doi:10.1002/aur.2777



99. St-Louis AC, Rapaport M, Chénard Poirier L, Vallerand RJ, Dandeneau S. On Emotion Regulation Strategies and Well-Being: The Role of Passion. *Journal of Happiness Studies*. 2021;22(4):1791-1818; doi:10.1007/s10902-020-00296-8

**Table 1.** Demographic characteristics ( $n = 108$ )

Variable	Full sample			
	<i>n</i>	%	M	SD
Age	-	-	23.66	5.11
<b>Gender</b>				
Female	57	52.7		
Male	43	39.8		
Other	7	6.5		
Missing	1	1		
<b>Occupation</b>				
Student	59	54.6		
Employed	18	16.7		
Unemployed	18	16.7		
Other	3	2.8		
Missing	10	9.2		
<b>Education</b>				
High School degree	15	13.9		
Collegiate or professional degree	26	24.1		
University degree	41	38.0		
No degree/currently in high school	26	24.0		
<b>Relationship status</b>				
Single	75	69.4		
Partner	33	30.6		
<b>Country</b>				
Canada	80	73.4		
France	27	25.7		
Belgium	1	0.9		

**Table 2.** Interest category and examples

Category (n; gender)	Activity examples	Quote	Passion levels
Gaming (n = 26; W = 6, M = 20)	Super Smash Bros	<i>"I play video games that involve strategy, which allows me to express my need to plan and manage things. Moreover, it is a passion that I can share with other gamers. It allows me to easily reach out and meet people, otherwise introducing myself to others in everyday life is difficult. With video gaming, I can get in touch with people that I know I have things in common with!"</i>	HP = 5.00 OP = 3.50 PC = 6.00
	Pokemon, Super Smash Bros	<i>"I enjoy playing with two video games. Either Pokemon or Super Smash Bros. I alternate between these two games in a cycle and never play other games. I typically change game every two months."</i>	HP = 3.83 OP = 4.50 PC = 6.80
Knowledge acquisition (n = 26; W = 16, M = 8, NB = 2)	Biology	<i>"This passion takes up most of my time during the day. I am lucky enough to study and work in this field that is my passion. My classes can be boring considering that I know most of what is being taught. To satisfy my curiosity, I constantly buy books on this topic, that I read as soon as I wake up in the morning."</i>	HP = 6.17 OP = 6.00 PC = 7.00
	Linguistics	<i>"I'm interested in learning about foreign languages. I particularly enjoy etymology and understanding the syntax of words. I read many grammar books and dictionaries and build comparative tables to work out differences between elements of two languages or more."</i>	HP = 6.33 OP = 5.33 PC = 7.00
Active arts (n = 22; W = 14, M = 8)	Drawing	<i>"Drawing. I draw whales and draw stories that I invent. I like tracing the outer lines and using watercolor to fill in the shapes. I draw in different booklets and on different paper types (textures and sizes). I especially enjoy finishing a drawing and start making a new one."</i>	HP = 5.33 OP = 5.83 PC = 6.00
	Singing	<i>"I have many interests, but singing is my favorite one. My echolalia is mostly present in form of singing and luckily, I am a soprano and can sing on pitch, which makes it easier for people around me."</i>	HP = 6.50 OP = 4.00 PC = 7.00

Television ( <i>n</i> = 11; <i>W</i> = 7, <i>M</i> = 2, <i>NB</i> = 1)*	Watching tv series	<i>“Watching anime series and movies”</i>	<i>HP</i> = 5.33 <i>OP</i> = 2.83 <i>PC</i> = 5.40
Reading ( <i>n</i> = 11; <i>W</i> = 8, <i>M</i> = 1, <i>NB</i> = 1)	Reading books	<i>“Reading. I read whatever I can find in different circumstances. For example, I enjoy reading books or any description, even posters when I walk on the sidewalks, and pamphlets in a waiting room.”</i>	<i>HP</i> = 5.67 <i>OP</i> = 6.83 <i>PC</i> = 7.00
Sports ( <i>n</i> = 6; <i>W</i> = 2, <i>M</i> = 3, <i>NB</i> = 1)	Cycling	<i>“I love cycling, more specifically road cycling, I like watching it and practicing the sport as well.”</i>	<i>HP</i> = 3.83 <i>OP</i> = 4.67 <i>PC</i> = 6.20
Animals ( <i>n</i> = 4; <i>W</i> = 2, <i>NB</i> = 2)	Taking care of pets	<i>“Taking care of my animals and their habitats. I own many animals (birds, cats, a dog, fishes, hedgehog, etc.), but my passion is taking care of my geckos and the insects surrounding them. I also take care of plants in their environments to create a rich ecosystem that benefits all species involved.”</i>	<i>HP</i> = 7.00 <i>OP</i> = 3.33 <i>PC</i> = 6.60
Passive arts ( <i>n</i> = 2, <i>W</i> = 2)	Listening to music	<i>“I listen to music with my earphones alone in my room. I walk around my room while listening to music. I can listen to the same song on repeat.”</i>	<i>HP</i> = 5.17 <i>OP</i> = 3.50 <i>PC</i> = 6.6

---

*Note.* *HP* = Harmonious passion; *M* = Men; *NB* = Non-binary; *PC* = Passion criteria; *OP* = Obsessive passion; *W* = Women. \*One participant with missing data for gender was in this category

**Table 3.** Descriptive statistics and correlations ( $n = 108$ )

Variable	n	M	SD	1	2	3	4	5	6	7	8	9	10	11	12	13
1. HP	108	5.41	1.09	-												
2. OP	108	4.31	1.44	-.13	-											
3. Flow	103	5.57	1.01	.26**	-.04	-										
4. Positive emotions	99	4.52	0.56	.32***	.13	.29**	-									
5. Negative emotions	99	1.69	0.66	-.35***	.21*	-.37***	-.10	-								
6. Conflict	107	3.82	1.68	-.14	.79***	-.08	.16	.19	-							
7. Rumination	98	4.02	1.68	.04	.70***	.04	.16	.24*	.59***	-						
8. Interpersonal relationships	92	3.85	0.87	.22*	-.22*	.17	.20	-.02	-.21*	-.14	-					
9. Well-being	90	3.72	1.8	.29**	-.31**	.11	.06	-.03	-.28**	-.22*	.36***	-				
10. Physical health	90	4.78	1.44	.08	-.38***	.01	-.08	-.08	-.29**	-.23*	.06	.46***	-			
11. Performance	89	4.05	1.71	.26*	-.47***	.22*	.09	-.15	-.40***	-.28**	.32**	.55***	.38***	-		
12. Contribution	89	5.17	1.53	.39***	-.24*	.14	.14	-.23*	-.16	-.15*	.23*	.29***	.26*	.55***	-	
13. OFIS	92	4.18	1.28	.35***	-.44***	.17	.09	-.16	-.36***	-.28**	.47***	.86***	.62***	.78***	.79***	-

Note.  $p < 0.1$ ; \* $p < .05$ ; \*\* $p < .01$ ; \*\*\* $p < .001$ .

**Figure legends**

**FIG. 1** Characterizing passion in autism. Each data point represents a participant and its respective levels of HP and OP ( $n = 108$ ).

**FIG. 2** Structural equation modelling results ( $n = 108$ ). Path coefficients are presented as standardized coefficients. † $p < 0.1$ ; \* $p < .05$ ; \*\* $p < .01$ ; \*\*\* $p < .001$